

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

AD-A247 038



due to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

DT DATE

3. REPORT TYPE AND DATES COVERED

ANNUAL 1 Nov 90 - 31 Oct 91

4. TITLE AND SUBTITLE

PROBING COSMIC INFRARED SOURCES: A COMPUTER MODELING APPROACH

5. FUNDING NUMBERS

GR - AFOSR-89-0104
PE - 61102F
PR - 2311
TA - A1

(2)

6. AUTHOR(S)

Dr Chun Ming Leung

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Physics Department
Rensselaer Polytechnic Institute
Troy, NY 12180-3590

8. PERFORMING ORGANIZATION REPORT NUMBER

AEOSR-TR- 92 0106

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Dr Henry R. Radoski
AFOSR/NL
Building 410
Bolling AFB DC 20332-6448

10. SPONSORING/MONITORING AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

DTIC
ELECTE
MAR 09 1992
S D D

12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release;
distribution unlimited

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

The research in infrared astronomy has modeled several classes of infrared sources and modified and generalized existing computer codes. The effects of temperature fluctuations due to small grains on the energy spectrum and infrared surface brightness of interstellar clouds heated externally by the interstellar radiation field were studied. A detailed study has begun on the grain formation problem in stellar outflows. The effects of fractal dust grains on the spectrum of infrared sources has been studied. Models are being developed of the physical conditions in the circumstellar envelopes of evolved stars. The chemistry of two dense interstellar clouds is being studied. The effect of temperature dependent opacity on the emergent spectra of infrared sources is being investigated.

14. SUBJECT TERMS

15. NUMBER OF PAGES

5

16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

19. SECURITY CLASSIFICATION OF ABSTRACT

UNCLASSIFIED

20. LIMITATION OF ABSTRACT

UNLIMITED

Annual Technical Report

PI: Chun Ming Leung
Institution: Rensselaer Polytechnic Institute
AFOSR-89-0104

From 01 Nov 90 to 31 Oct 91

Received: 13 December 1991

Accepted: 16 December 1991

Narrative:

The research in infrared astronomy has modeled several classes of infrared sources and modified and generalized existing computer codes. The effects of temperature fluctuations due to small grains on the energy spectrum and infrared surface brightness of interstellar clouds heated externally by the interstellar radiation field were studied. A detailed study has begun on the grain formation problem in stellar outflows. The effects of fractal dust grains on the spectrum of infrared sources has been studied. Models are being developed of the physical conditions in the circumstellar envelopes of evolved stars. The chemistry of two dense interstellar clouds is being studied. The effect of temperature dependent opacity on the emergent spectra of infrared sources is being investigated.

PM: H.R.Radoski

Accession For	
NTIS CRA&I	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability	
Dist	Avail Spec
A-1	



92-05706

92 3 03 185



Rensselaer

December 6, 1991

Department of Physics

Dr. Henry R. Radoski, Director
Space Physics Program
Life Sciences Division
Air Force Office of Scientific Research
AFOSR/NP, Building 410
Bolling Air Force Base, D. C. 20332-6448

Re.: 1990-91 Research Progress Report (Grant AFOSR-89-0104)

Dear Dr. Radoski:

Thank you for your recent approval of a six-month no-cost extension of my AFOSR grant supporting the project: *"Probing Cosmic Infrared Sources: a Computer Modeling Approach"*. As you suggested, in lieu of a lengthy third annual technical report, I am submitting a brief summary of the research activities in the past twelve months. A detailed final technical report will be submitted after the expiration of this grant on April 30, 1992, as required.

Personnel:

At present the research group working on this project consists of 6 people: Chun M. Leung (PI), Michael P. Egan, Steven D. Doty, Michael E. Fogel, Monika E. Kress and Curtis Potterveld (all graduate students). Both Egan and Doty are Ph. D. candidates. The two postdoctoral fellows, Drs. Steven B. Charnley and K. N. Nagendra, completed their terms of appointments and left to take up positions elsewhere: Charnley is now at the NASA Ames Research Center while Nagendra is at the Indian Institute of Astrophysics at Bangalore in India. In addition, three undergraduates were involved in research this past summer.

Research:

The research effort has gone into modifying and generalizing existing computer codes as well as in the modeling of several classes of infrared sources.

- (1) We continued to study the effects of temperature fluctuations due to small grains on the energy spectrum and infrared surface brightness of interstellar clouds heated externally by the interstellar radiation field. Radiation transport models with conventional large grains, very small grains, and polycyclic aromatic hydrocarbons (PAHs) are constructed. Applying the model results to the IRAS observations of the diffuse cloud in Chamaeleon, it is found that the 25 μm emission is most likely due to small grains which also contribute significantly to the observed 12 μm emission. However, the majority of the 12 μm flux is due to PAHs. The emission at 60 and 100 μm is mostly due to large grains (with radius $\sim 0.1 \mu\text{m}$) heated under equilibrium conditions. Furthermore, to produce the observed limb-brightening at various infrared wavelengths, the spatial distribution of the small grains and PAHs must be more extended than the large grains.
- (2) We have begun a detailed study of the grain formation problem in stellar outflows. Grain formation consists of nucleation followed by growth. Most studies have used classical nucleation theory to describe the formation rate of grain nuclei in a supersaturated vapor, under the assumption of local thermodynamic equilibrium. However, the nucleation process is best determined by solving a system of kinetic

equations. The growth of the nuclei is then determined by solving a set of moment equations which are highly dependent on the nucleation rate of particles in the gas. To test the validity of classical nucleation theory in astrophysical environments, a truncated set of kinetic equations is solved simultaneously with the moment equations, thus allowing a self-consistent treatment of grain nucleation and growth. It is found that classical nucleation theory allows nucleation of grains to occur at a much lower supersaturation than is calculated from the kinetic equations, thus predicting dust formation too near a star, overestimating the number of dust grains produced, and underestimating the average grain size. The effect of these results on radiative transfer in the dust shell is being investigated.

- (3) We have studied the effects of fractal dust grains on the spectrum of infrared sources by considering non-spherical dust grains produced by two fractal growth processes: particle-by-particle aggregation or cluster-by-cluster aggregation. By approximating a fractal grain as a collection of dipoles, the discrete dipole approximation is used to calculate the grain opacity. Optical constants for both bulk material and amorphous clusters are utilized. Using these dust opacities for fractal grains, radiation transport models of infrared sources are constructed to study the effect of grain size, shape, and composition on the IR spectrum. It is found that, compared to models with spherical grains of the same composition and volume, models with fractal carbon grains show a shift in the peak flux toward longer wavelengths. The less compact fractal dust grains tend to be cooler than spherical grains. These differences are due to a higher ratio of geometrical cross section to volume for fractal grains. For fractal grains with the same ratio the overall grain shape plays only a minor role.
- (4) We have begun a study to model in detail the physics and chemistry of the circumstellar envelopes of evolved stars. The goal is to develop a comprehensive model in which both chemical evolution and radiation hydrodynamics are treated self-consistently. As an example, the effect of shielding of the ambient interstellar radiation by dust and self-shielding of H_2 and CO on the photoionization rates are considered by solving the radiation transport problem accurately. Model results will be compared with available observations of evolved stars.
- (5) We have begun a study to understand the chemistry of the two dense interstellar clouds, L134N and TMC-1. These clouds have similar physical conditions but display remarkable differences in chemical abundances. In particular, whereas TMC-1 exhibits relatively large abundances of many complex, unsaturated, carbon-rich species, L134N appears to have larger abundances of some oxygen-rich species as well as saturated simple molecules. This discrepancy may be due to a difference in the initial gas-phase $[C]/[O]$ ratio or due to the clouds being in different stages of evolution. To delineate the effects of different initial composition versus cloud evolution, the chemical model of Herbst and Leung are used to study the chemical evolution of these two clouds. In the models the original reaction network has been expanded to include 3517 reactions involving 385 species. It is found that certain molecules may prove to be particularly sensitive probes of the initial $[C]/[O]$ ratio and/or the evolutionary phase of the cloud. In particular, the ratio $[NO]/[CO]$ can be used to differentiate between whether an initial high $[C]/[O]$ ratio (relative to the cosmic value) is due to carbon enrichment or to oxygen depletion, e.g., $[NO]/[CO] > 1$ for carbon enrichment while $[NO]/[CO] < 1$ for oxygen depletion.
- (6) We have begun to study the effect of temperature-dependent opacity on the emergent spectra of infrared sources. Laboratory evidence indicates that optical constants of grain material depend on the grain temperature. Radiation transport models are constructed in which the dust opacity is allowed to vary with grain temperature. It is found that a temperature-dependent dust opacity affects both the equilibrium grain temperature and the emergent spectrum. The effect is most pronounced on the detailed shape of spectral features. Models are being constructed to study the effects on the intensity and shape of the $10\ \mu m$ and $20\ \mu m$ features of silicate grains.

Preliminary results of the research in (2), (3), and (5) will be presented as three poster papers in the upcoming meeting of the American Astronomical Society in Atlanta.

I greatly appreciate the support by AFOSR of this research program. I hope you have received my recent proposal entitled "*Computer Modeling Studies of the Infrared Celestial Backgrounds*" for the DoD program of Augmentation Awards for Science and Engineering Research Training and that the proposal will receive favorable consideration.

Sincerely,

A handwritten signature in cursive script, reading "Chun Ming Leung".

Chun Ming Leung
Professor of Physics